


Latinos and HIV/AIDS

December 2006

Latinos in the United States continue to be affected by the HIV/AIDS epidemic, accounting for a greater proportion of AIDS cases than their representation in the U.S. population overall, and the second highest AIDS case rate in the nation, by race/ethnicity.¹ The epidemic has had a disproportionate impact on Latinas and young adults, and the impact of HIV/AIDS among Latinos varies across the country and by place of birth.^{1,2} Moreover, studies have shown that Latinos with HIV/AIDS may face additional barriers to accessing care than their white counterparts.^{3,4,5} Today, there are approximately 1.2 million people living with HIV/AIDS in the U.S., including about 200,000 Latinos.⁶ As the largest and fastest growing ethnic minority group in the U.S., addressing the impact of HIV/AIDS in the Latino community takes on increased importance in efforts to improve the nation's health.

- The number of Latinos living with AIDS has also increased over time, in part due to treatment advances but also to the epidemic's continued impact. Estimated AIDS prevalence among Latinos increased by 33% between 2001 and 2005, compared to a 21% increase among whites.¹
- The number of deaths among Latinos with AIDS remained stable between 2001 and 2005, while both Blacks and whites experienced slight decreases.¹

Figure 1: Estimated AIDS Diagnoses & U.S. Population by Race/Ethnicity, 2005^{1,7,8}




Snapshot of the Epidemic

- Although Latinos represent approximately 14% of the U.S. population⁸, they account for 19% of the AIDS cases diagnosed in 2005 and 16% of the AIDS cases diagnosed since the start of the epidemic (Figure 1).^{1,7} Latinos account for 18% of HIV/AIDS cases diagnosed in 2005 in the 33 states with confidential names-based reporting.^{1,7}
- The AIDS case rate per 100,000 among Latino adults/adolescents was the second highest of any racial/ethnic group in the U.S. in 2005—3.5 times that of whites, but about one-third that of Blacks (Figure 2).^{1,9}
- HIV was the 6th leading cause of death for Latinos aged 25-34 in 2002, the same ranking as for whites. HIV was the 3rd leading cause of death for Blacks in this age group.¹⁰ In 2003, HIV deaths rates per 100,000 population, aged 25-44, were higher among Latinos (10.3 for men and 3.8 for Latinas) compared to whites, although they were highest for Blacks.¹¹

Key Trends and Current Cases

- Latinos account for a growing share of AIDS diagnoses over time, rising from 15% in 1985 to 19% in 2005; in recent years, this share has remained relatively stable.^{1,7,12}

Figure 2: AIDS Case Rate per 100,000 Population by Race/Ethnicity for Adults/Adolescents, 2005^{1,9}



Women and Young People

- Among women, Latinas account for 16% of new AIDS cases in 2005; Black women account for 67% and white women account for 16%.^{1,7,9}
- Latinas represent 22% of AIDS cases diagnosed among Latinas in 2005; by comparison, white women represent 14% of cases among whites, and Black women represent 35% of cases diagnosed among Blacks.^{1,9}
- The AIDS case rate per 100,000 among Latinas (26.4) was nearly 6 times higher than the case rate for white women (2.1).^{1,9}
- Latino teens, aged 13-19, accounted for 14% of AIDS cases among teens compared to 16% of all U.S. teens in 2004.² Latinos aged 20-24 accounted for 23% of new AIDS cases reported among young adults, but represented 18% of U.S. young adults, in 2004.²


Transmission

- HIV transmission patterns among Latino men vary from those of white men. Although both groups are most likely to be infected through sex with other men, white men are more likely to have been infected this way. Heterosexual transmission and injection drug use account for a greater share of infections among Latino men than white men.^{1,13}
- Latinas are somewhat more likely to have been infected through heterosexual transmission than white women, although this is the most common transmission route for both groups and for women overall. White women are somewhat more likely to have been infected through injection drug use than Latinas.^{1,13}

- Studies have found high HIV/AIDS prevalence among Latino men who have sex with men (MSM).¹⁴ A study in 5 major U.S. cities found that 17% of Latino MSM in the study were infected with HIV. Prevalence among white MSM was 21% and among Black MSM, 46%, the highest of any group.¹⁵ Knowledge of HIV status among those already infected was also very low.¹⁴

- Insurance status also varies at the time of HIV diagnosis. Analysis of data from 25 states between 1994 and 2000 found that Latinos were less likely than whites to have private coverage and more likely to be covered by Medicaid at the time of their HIV diagnosis. A third of Latinos were uninsured at the time of their diagnosis, higher than other groups.²⁰

Figure 3: Number of Latinos Estimated to be Living with AIDS: Top 10 States/Areas, 2005¹⁶



Geography

Although AIDS cases among Latinos have been reported throughout the country, the impact of the epidemic is not uniformly distributed:

- AIDS case rates per 100,000 among Latinos are highest in the eastern part of the U.S., particularly in the Northeast.¹⁶ The Northeast also has the greatest proportion of Latinos estimated to be living with AIDS (37% in 2005) and new AIDS cases among Latinos (33%).^{16,17}
- AIDS prevalence among Latinos is clustered in a handful of states, with 10 states accounting for 89% of Latinos estimated to be living with AIDS in 2005. New York, California, and Puerto Rico top the list (Figure 3). Ten states also account for the majority of newly reported AIDS cases among Latinos (86% in 2005).^{16,17}
- AIDS cases among Latinos vary by place of birth. Latinos born in the U.S. accounted for 41% of estimated AIDS cases among Latinos in 2005, followed by Latinos born in Puerto Rico (22%) and Mexico (22%).^{1,18} HIV transmission patterns among Latinos also vary by place of birth.¹

Access to and Use of the Health Care System

The HIV Cost and Services Utilization Study (HCSUS), the only nationally representative study of people with HIV/AIDS receiving regular or ongoing medical care for HIV infection, found that Latinos fared more poorly on several important measures of access and quality, differences that diminished over time but were not completely eliminated.³ In addition, HCSUS found that Latinos were more likely to report postponing medical care due to factors such as lack of transportation.⁴ Latinos were also more likely than whites to delay care after an HIV diagnosis.⁵

Health Insurance

Having health insurance, either public or private, improves access to care. Insurance coverage of those with HIV/AIDS varies by race/ethnicity, as it does for the U.S. population overall.

- The HCSUS study found that Latinos with HIV/AIDS were more likely to be publicly insured or uninsured than their white counterparts, with half relying on Medicaid compared to 32% of whites. Approximately one quarter of Latinos with HIV/AIDS (24%) were uninsured compared to 17% of whites. Latinos were also about half as likely to be privately insured than whites (23% compared to 44%).¹⁹

HIV Testing

- Among the U.S. population overall, Latinos are more likely than whites to report ever having ever been tested for HIV (54% compared to 45%). However, these self-reported testing rates may be overestimates, since 20% of Latinos assumed that the test was a routine part of an exam.²¹
- Among those who are HIV positive, CDC data indicate that more than one in four Latinos (43%) were tested for HIV late in their illness—that is, diagnosed with AIDS within one year of testing positive (in 33 areas with HIV reporting); by comparison, 40% of Blacks and 37% of whites were tested late.¹

Concern About HIV/AIDS²¹

- A recent survey found that Latinos express concern about HIV/AIDS. Nearly a quarter of Latinos named it as the most urgent health problem facing the nation, ranked second after cancer. More Latinos believe the U.S. is making progress on the domestic epidemic (39%) than losing ground (30%), as do whites; by contrast, Black Americans are more likely to say the U.S. is losing ground.
- Almost half (46%) of Latinos say they think AIDS is a more urgent problem in their community than it was a few years ago compared to 15% of whites. Although 31% of Latinos say they are personally very concerned about becoming infected with HIV, this proportion has declined since the mid-1990's.

References

- 1 CDC, *HIV/AIDS Surveillance Report*, Vol.17; 2006.
- 2 CDC, *HIV/AIDS Surveillance in Adolescents and Young Adults* (through 2004).
- 3 Shapiro MF et al., "Variations in the Care of HIV-Infected Adults in the United States." *JAMA*, Vol. 281, No.24; 1999.
- 4 Cunningham WE et al., "The Impact of Competing Subsistence Needs and Barriers to Access to Medical Care for Persons with Human Immunodeficiency Virus Receiving Care in the United States." *Medical Care*, Vol.37, No.12; 1999.
- 5 Turner BJ et al., "Delayed Medical Care After Diagnosis in a U.S. Probability Sample of Persons Infected with the Human Immunodeficiency Virus." *Archives of Internal Medicine*, Vol.160; 2000.
- 6 Kaiser Family Foundation calculations based on: Glynn MK and Rhodes P, "Estimated HIV Prevalence in the United States at the end of 2003." Presentation, National HIV Prevention Conference; June 2005.
- 7 Calculations based only on cases for which race/ethnicity data were provided.
- 8 U.S. Census Bureau, 2005 Population Estimates.
- 9 Includes reported cases among those 13 years of age and older. Estimates do not include cases from the U.S. dependencies, possessions, and associated nations, and cases of unknown residence.
- 10 NCHS, "Deaths: Leading Causes for 2002," *NVSR*, Vol. 53, No. 17; 2005.
- 11 NCHS, *Health, United States, 2005*.
- 12 CDC data request; January 2006.
- 13 CDC, *HIV/AIDS Surveillance by Race/Ethnicity* (through 2004).
- 14 CDC, *Fact Sheet: HIV/AIDS Among Men Who Have Sex with Men*, July 2005.
- 15 CDC, "HIV Prevalence, Unrecognized Infection, and HIV Testing Among Men Who Have Sex with Men – Five U.S. Cities, June 2004–April 2005," *MMWR*, Vol. 54, No. 24; June 24, 2005.
- 16 Kaiser Family Foundation, www.statehealthfacts.org. Data Source: Centers for Disease Control and Prevention, Division of HIV/AIDS Prevention-Surveillance and Epidemiology, Special Data Request; November 2006.
- 17 Estimates include U.S. dependencies, possessions, and associated nations, and cases of unknown residence.
- 18 Calculations based only on cases for which data by place of birth were provided.
- 19 Fleishman JA. Personal Communication, Analysis of HCSUS Data; January 2002.
- 20 Kaiser Family Foundation analysis of CDC data.
- 21 KFF, *Survey of Americans on HIV/AIDS*; 2006

The Kaiser Family Foundation is a non-profit, private operating foundation dedicated to providing information and analysis on health care issues to policymakers, the media, the health care community, and the general public. The Foundation is not associated with Kaiser Permanente or Kaiser Industries.
Additional copies of this publication (#6007-04) are available on the Kaiser Family Foundation's website at www.kff.org.